



KNOMAD WORKING PAPER 4

What really is Brain Drain? Location of Birth, Education, and Migration Dynamics of African Doctors

Çağlar Özden
David Phillips

February 2015



Federal Ministry
for Economic Cooperation
and Development



Centre for International
Migration and Development
a joint operation of GIZ and the
German Federal Employment Agency



REGERINGSKANSLIET

Government Offices
of Sweden



Schweizerische Eidgenossenschaft
Confédération suisse
Confederazione Svizzera
Confederaziun svizra

Swiss Agency for Development
and Cooperation SDC



WORLD BANK GROUP

The KNOMAD Working Paper Series disseminates work in progress under the Global Knowledge Partnership on Migration and Development (KNOMAD). A global hub of knowledge and policy expertise on migration and development, KNOMAD aims to create and synthesize multidisciplinary knowledge and evidence; generate a menu of policy options for migration policy makers; and provide technical assistance and capacity building for pilot projects, evaluation of policies, and data collection.

KNOMAD is supported by a multi-donor trust fund established by the World Bank. Germany's Federal Ministry of Economic Cooperation and Development (BMZ), Sweden's Ministry of Justice, Migration and Asylum Policy, and the Swiss Agency for Development and Cooperation (SDC) are the contributors to the trust fund.

The views expressed in this paper do not represent the views of the World Bank or the sponsoring organizations.

All queries should be addressed to KNOMAD@worldbank.org. KNOMAD working papers and a host of other resources on migration are available at www.KNOMAD.org.

What really is Brain Drain?

Location of Birth, Education, and Migration Dynamics of African Doctors*

Çaglar Özden and David Phillips†

Abstract

Proper analysis of skilled migration flows and their impact requires joint identification of where migrants were born, educated, and when they moved to the destination country. To highlight these issues and identify key patterns in career paths, we focus here on doctors practicing in the United States who were born and/or trained in Africa. We overcome data constraints by merging data from the American Medical Association (AMA) and American Community Survey (ACS) via propensity score matching techniques. Our results show that the standard assumptions on skilled migration lead to considerable overestimation of its extent and hide several economically important migration patterns. We find that almost half of African-born doctors were trained outside their country of birth. On the flip side, around 15 percent of all doctors trained in Africa were actually born outside the continent. There is significant variation across countries in terms of age of migration levels, implying that many African doctors migrate after years of service and that their human capital is not completely lost to their birth countries. In short, global labor and education markets for high-skilled professionals are integrated in more nuanced and thought-provoking ways than assumed in the literature.

Keywords: Brain drain, high-skilled migration, doctors, Africa, United States

* Paper produced under KNOMAD's Thematic Working Group (TWG) on Skilled Migration. KNOMAD is headed by Dilip Ratha, the TWG on Skilled Migration is chaired by Çaglar Özden, and the focal point is Kirsten Schuettler. The authors thank the American Medical Association for the data used in this paper. The authors are grateful to Erhan Artuc, Michel Beine, Simone Bertoli, Michael Clemens, Frederic Docquier, Gordon Hanson, Robert Lucas, Manjula Luthria, Aaditya Mattoo, David McKenzie, Hillel Rapoport, Andrew Viteritti, Mathis Wagner, and seminar participants at Georgetown University, the World Bank, CEPR Conference at Nottingham University, the 7th International Conference on Migration and Development, Syracuse University, and Western Michigan University for comments and suggestions. This paper also reflects comments received through the KNOMAD peer review process.

† Çaglar Özden is Lead Economist at the World Bank and David Phillips is Assistant Professor of Economics at Hope College, Michigan. The authors may be contacted at cozden@worldbank.org and phillipsd@hope.edu.

Table of Contents

1. Introduction.....	1
2. Data	4
2.1. American Medical Association Data	4
2.2. American Community Survey Data	6
3. Estimation Methods	7
3.1. Trained in Africa	8
3.2. Trained Outside Africa.....	9
3.3. Estimating Age at Migration.....	12
3.3.1. Estimating Age at Migration for Those Born and Trained in Africa.....	13
3.3.2. Estimating Age at Migration for Those Trained Outside Africa.....	14
3.3.3. Combining Age at Migration Distributions	14
4. Results	14
4.1. Propensity Score Matching Results.....	14
4.2. Missing Location of Birth and Education in the Databases.....	15
4.3. Patterns in Location of Birth and Education	16
4.4. Age of migration.....	20
5. Conclusion	24
References	26

1. Introduction

High-skilled migration from poorer to wealthier countries, commonly referred to as “brain drain”, is viewed with significant unease due to the vital role played by human capital in economic growth and development processes. The key questions that link the migration and growth/development literatures¹ concern where human capital is created and where it is utilized. The literature makes several simplifying assumptions, such as education being obtained in the birth country, to deal with the data constraints that emerge in answering these questions. We follow a different route and develop an empirical methodology that enables us to *jointly* identify where the migrant doctors are born, where they are educated and at what age they migrate, using African-born and/or -educated doctors in the United States as an example. Our results show that the standard assumptions lead to considerable overestimation of the extent of high-skilled migration and hide several economically important migration patterns.

Over the last decade, rigorous empirical research has begun to catch up with the earlier analytical models² and aid policy evaluation (Hanson 2010; Docquier and Rapoport 2012). Newly available migration databases and the subsequent economic analyses depend on two critical assumptions. First, country of birth is generally equated with the country of education.³ This definition results from reliance on census-type data sources where birth location is used as the main determinant of migration status. As a result, human capital obtained in the destination country is generally attributed to the birth country. Second, time spent in the country of education after graduation is almost always ignored. The literature often assumes that a migrant moves right after graduation and that his human capital is only utilized in the destination country. Since most skilled migration takes place from poor developing countries to wealthier OECD countries, both of these assumptions are likely to lead to biases regarding its extent and impact. However, an Ethiopian-born, British-educated professional who moved to the United States in the middle of his career matters for the education, labor and fiscal policies of all three countries. We need more detailed information on the educational and professional paths to properly identify skilled migration patterns and their effect on all of the countries that these paths happen to cross.

Collection of extensive personal data through surveys is one option to address these issues, but likely to be prohibitively costly. Instead, we develop a propensity score matching technique to merge two different existing data sets with complementary types of information. The first is the administrative data from the American Medical Association (AMA) that covers education histories of doctors. The second source is the census-type data from the American Community Survey (ACS) on the place of birth and age of migration of a representative sample of African-born doctors. Our matching technique links doctors that are similar in the two datasets to fully characterize their career paths. We are freed from making

¹ We interpret “growth” literature broadly to include a wide range of issues in development, labor market, and education.

² Bhagwati and Hamada (1974) and Stark, Helmenstein, and Prskawetz (1997) are examples of earlier theoretical work.

³ See Docquier and Marfouk (2006), Özden et. al. (2011), and Artuc et.al. (2015), as examples.

the two assumptions listed above and are able to identify where our African doctors were born, where they were educated, and at what age they migrated to the United States.

Our results identify complex migration patterns and detailed answers to the questions on where human capital is created and utilized. For example, almost half of the African-born doctors were not trained in Africa but in many different parts of the world, including South Asia and the Caribbean. These patterns imply many existing studies might be overestimating brain drain levels and certain effects, such as the fiscal burden on countries of birth. On the other hand, 15 percent of the African-trained doctors came from countries outside Africa, including the West, challenging the widely shared belief that skilled migration is a one-way drain from poorer to wealthier countries. In terms of when they migrate, we find that many doctors spend quite a few years between their graduation at home and migration to the United States, implying that countries of education do not immediately and fully lose the human capital generated within their borders. Finally, we observe substantial heterogeneity across African countries in terms of all of our results. This should caution all policymakers against making generalizations based on evidence from a single country.⁴

There are several reasons why we focus on doctors, especially those from Africa.⁵ The first is data availability. Due to licensing requirements in medical professions, administrative data records on medical training (school name and location) can be obtained for virtually every doctor in the United States. No other profession has such requirements and extensive data availability. Second, migration of health professionals holds a prominent place in the literature on skilled migration. There are few issues that generate as much emotion and controversy as the impact of medical migration, especially of African doctors to OECD countries (see, for example, Chen and Boufford 2005). The concern is that the exodus of African doctors will undermine the quality and delivery of healthcare services with long-lasting negative effects as many African countries struggle with crushing healthcare burdens. As a result, migration of African doctors has become the symbol of brain drain in many policy debates. Former WHO Director General Lee Jong-wook's (2006) statement that migration of African doctors leads to "loss of hope and years of investment" succinctly captures the existing sentiment.

The extent of this "loss", however, depends crucially on where education is obtained. This is especially true in the case of medical professions where education is expensive and publicly financed out of rather tight budgets, demand is outstripping supply in almost every country and provision of health services generates important development spillovers. As illustration, consider a radiologist who was born in Ghana, came to the United States to complete his tertiary and medical education, and continues to practice there – as opposed to a surgeon who came to the United States after completing her medical school in Ghana. By the country of birth criteria, both are part of Ghanaian medical brain drain. However, arguments against migration due to the fiscal burden imposed (e.g. Bhagwati and Hamada 1974) would imply that our surgeon creates more of a "loss" for Ghana since she was trained at home while the radiologist did not impose any financial or opportunity costs since he was trained overseas. On the other hand, someone accounting for the effect of migration on overall skill levels, health care

⁴ See Gibson and McKenzie, 2011 for a detailed discussion on brain drain.

⁵ See Easterly and Nyarko (2009) for an interesting discussion on skilled migration from Africa.

staffing, or health outcomes in Ghana may want to use a country of birth definition, because a high-ability student migrating at an early age could lead to reduced positive externalities. The situation becomes further complicated if we extend our list to include a pediatrician who was born in Ghana, trained in Canada, and migrated to the United States. For which, exactly, does she represent a “loss”? Canada or Ghana? A final issue arises when we focus on the age at which our Ghanaian-trained surgeon migrates to the United States. In terms of human capital mobility and impact on Ghana, migrating immediately after medical school differs significantly from migrating at age 45 after two decades of practice in Ghana; the latter presumably generates social externalities that compensate for the public funds that paid for her education. The Ministry of Health and the University of Ghana Medical School would need this information in order to better design their long-term human resource and education policies. One of our goals is to highlight the importance of the differences between our Ghanaian radiologist, surgeon, and pediatrician with respect to public policy. These differences are not captured and analyzed in the existing literature but matter greatly for welfare analysis and policy discussion about origin, transit, and destination countries.

Data constraints are the main obstacles in the high-skilled migration literature, especially in answering questions regarding location of education and timing of migration. The last decade witnessed an impressive leap forward but there are many remaining challenges, as discussed in detail by Hanson (2010). Among the earlier work, the Docquier-Marfouk (2008) and OECD-DIOC bilateral skilled-migration databases include only OECD destinations and use a location of birth classification. These were extended by OECD DIOC-E to cover numerous non-OECD destinations and predict skilled-migration levels for the whole world (Artuc et al. 2015). The shortcomings created by the location of birth classification have received significant attention (Docquier and Rapoport 2012). Realizing this dilemma, Beine, Docquier, and Rapoport (2007) and OECD extended their original database, dividing bilateral migration stocks into several sub-categories based on age-of-migration cutoffs as a proxy for location of education. This is a much-needed advance in terms of the data but can still leave unresolved issues. For example, we can classify tertiary-educated migrants as birth (destination) country-educated if they arrived after (before) age 22, presumably the age at which most people complete their university education. However, age of university graduation varies widely across countries and professions, as we see in our data. Furthermore, it is possible that many migrants were trained in a country other than their birth or destination country, as confirmed by many African migrant doctors trained in Europe, South Asia, and the Caribbean.

Despite its importance, the literature on migration of healthcare professionals also ignores joint identification of location of birth and training. Most existing studies prefer to choose one criterion or the other, highlighting their reasons for the choice. For example, Clemens and Petterson (2008) use country of birth; Mullan (2005) and Okeke (2009) use country of training, and the most ambitious cross-country panels use a pragmatic mixture of the two depending on which is available from different destination countries (Bhargava and Docquier 2008; Bhargava, Docquier, and Moullan 2011). The empirical literature then uses these varied definitions to test common hypotheses regarding the determinants of medical migration, the possibility of medical brain gain, and potential health effects of physician migration using cross-country data (see above citations; Clemens 2007; Kangasniemi et al. 2007).

Popular press and even policy analyses provide a relatively simplistic view of skilled migration, in which a person is born and trained in a developing country and then migrates after graduation to a wealthier country where incomes are higher. This story is undoubtedly true for many migrants. However, examining the data on African-born and/or -trained doctors in the US indicates that this view oversimplifies reality. The lives and careers of African doctors, and probably other groups of skilled migrants, follow many paths that all carry unique public policy implications. We focus on African doctors due to availability of detailed education data and their prominent role in the brain drain debate. However, it is important to note that our methods and conclusions can probably be applied to other professions and regions. Further analysis will surely uncover other interesting patterns in the global labor market for skills. Completing this picture remains the critical issue.

The next section explains the data used in the paper, followed by a detailed description of the empirical methodology. We then present our main results, discussing the major distinctions between location of birth and education numbers for various countries in Africa. We follow up with the age of migration results and then conclude the paper.

2. Data

2.1. American Medical Association Data

Records from the American Medical Association's (AMA) Physician Masterfile are our main data source. The Masterfile provides academic, professional and basic demographic information on all doctors who have either entered a medical school in the United States or who were trained outside but obtained licenses in the United States. Since the Masterfile covers all points of entry to the medical profession in the United States, the database provides a comprehensive source of data for licensed doctors in the United States.⁶ In the data collection, the AMA coordinates with many agencies, institutions, and organizations, including medical schools, post-graduate programs, state licensing boards, the Educational Commission for Foreign Medical Graduates, and the American Board of Medical Specialties.

We obtained an extract from the Masterfile in 2011 including all doctors for whom the AMA lists (i) an African country as their country of birth or (ii) a medical school located in Africa as their location of training. The list includes all countries in sub-Saharan Africa as well North Africa. From this extract, we limit our focus to those doctors who are under the age of 65. This prevents counting retired doctors (who may still be listed in the Masterfile). In addition, this cutoff assists in matching AMA data to the American Community Survey (ACS) data that we describe later in this section. The individual records from the Masterfile include name, gender, age, name of medical school attended, year of graduation, year of birth, current address, and primary specialty.

⁶ See <http://www.ama-assn.org/ama/pub/about-ama/physician-data-resources/physician-masterfile.page> for a more detailed description of the data.

Because training information is required for accreditation and professional licensing, the data for school of medical training is complete. We identify the geographic location of each medical school listed using the International Medical Education Directory.⁷ For ease of presentation and due to the data requirements of our methods (see Section 3), we group all possible countries of training into 19 regions, of which 13 are in Africa and six outside the continent. Of the 13 African regions, 10 are relatively large individual countries: Egypt, Ethiopia, Ghana, Kenya, Liberia, Nigeria, South Africa, Sudan, Uganda, and Zimbabwe. The other three are geographic regions that capture the rest of the African countries, since these generally have small numbers of doctors in the United States. These regions are (i) Other East Africa, (ii) Other North Africa, (iii) Other West, Central, and Southern Africa. Finally, the six regions outside Africa are: (i) United States, (ii) Other English-speaking developed countries (United Kingdom, Canada, Australia, New Zealand, and Ireland), (iii) non-English speaking Europe, (iv) the Caribbean, (v) South Asia, and (vi) the Rest of the world. In this paper we refer to these 19 groupings broadly as countries, locations, and regions.

Summary statistics of the AMA Masterfile are presented in Table 1 (see end of document for all tables). Column 1 shows the location of training. In this context, Egypt leads the African countries with 27 percent of the total (of 15,191 doctors in the Masterfile) followed by Nigeria with 21 percent and South Africa with 9 percent. We should note that there are also many African-born doctors trained in the United States and they account for 17 percent of the sample. In total, 26 percent of the AMA file is composed of African-born doctors who were trained outside Africa. Column 3 lists the current mean age and Column 4 presents the fraction that is female for each location of training.

Location of birth is a key variable in the migration literature, but, unfortunately, it is not universally reported in the administrative AMA data. Non-response is quite common, with only 55 percent of doctors in the sample answering this question even though the AMA strongly encourages it. For instance, 63 percent of the African-trained doctors and 47 percent of the total observations do not report their country of birth.⁸ Finally, for those who do report country of birth, we group the locations in the same manner as we do locations of training. As shown in Column 2, of the total number of observations in the Masterfile, 17 percent were born in Egypt, 6 percent were born in Nigeria, and 5 percent in each of Ghana and Kenya.

At various points, we make use of other data from the AMA Masterfile including current age, age at graduation from medical school, and the census division in which the doctor practices. Gender and whether the doctor is a DO or an MD are also listed. We group listed medical specialties into 20 broad

⁷ The International Medical Education Directory is a public database of worldwide medical schools and is published as a joint collaboration of the Educational Commission for Foreign Medical Graduates and the Foundation for Advancement of International Medical Education and Research.

⁸ Non-reporting is particularly high among Nigerian-trained doctors under the age of 45. Because response rates are so low for this group, we code all such doctors as having a missing country of birth. There are 204 such observations out of 3,254 Nigerian-trained doctors. Due to unusually high reporting of the United States as a country of birth among the few young Nigerians who do report country of birth, dropping these observations generates conservative estimates of the number of African-trained but not African-born doctors.

specialty groups.⁹ We also assign a variable for whether the doctor has an ethnically Asian name. The fraction of the doctors with an Asian name is particularly high for Kenya, Uganda, and other East Africa, reflecting the historical ethnic composition of these countries due to their colonial legacies. This variable is useful because African doctors of South Asian descent may behave differently than other doctors in terms of their migration patterns.¹⁰ We assign this variable if the individual's last name occurred at least 100 times in the 2000 US Census and if at least 70 percent of individuals with this name reported their ethnicity as Asian.¹¹ Visual inspection confirms that these names are almost entirely not only Asian, but also South Asian specifically.

2.2. American Community Survey Data

The second data source is the weighted five-year sample of the American Community Survey (ACS) 2010 from the IPUMS. Our sample includes those individuals who report their birthplace to be in Africa and physician as their current occupation. As with the AMA data, we limit the sample to those under the age of 65. Most importantly for our purposes, the ACS data has complete information on country of birth as well as additional data from which we can compute age at migration. On the other hand, the ACS has two shortcomings. First, there is no information on training locations. Second, the available data only cover a sample of the African-born doctors, as opposed to the AMA Masterfile's complete universe of doctors.

We classify locations of birth into the same 13 groupings in Africa. Since the location of training is not available, we cannot identify any doctor who was born outside but trained in Africa in the ACS. This limitation forces us to use only the African-born observations from the sample. A small number of observations (36 out of 870) are identified only by region (for instance "East Africa, not stated/not elsewhere classified"). We classify these observations in the various "other" categories, according to the region. An additional 30 observations are only listed as born in Africa. Our treatment of these observations is described in the methods section. We also use current age and gender from the ACS data. We can calculate age at migration as the difference between current age and how many years the person has lived in the United States. Finally, we use state of residence to identify the census division in which the person resides.

Table 2 provides the basic summary statistics for the ACS data in which there are 17,940 African-born doctors. Of these, 23 percent were born in Egypt, 21 percent in Nigeria and 15 percent in South Africa, with the rest of Africa accounting for the remaining 41 percent. Column 3 presents the mean age, column 4 the fraction who are female, and the last column the mean age of migration to the United States. Egyptian and Sudanese doctors have the lowest ratio of females while Kenyans, Zimbabweans, and Ghanaians have the lowest average age of migration.

⁹ They are anesthesiology, cardiovascular, emergency, family, gastroenterology, infectious disease, internal, nephrology, neurology, OB/GYN, oncology, ophthalmology, orthopedics, pathology, pediatrics, psychiatry, pulmonary, radiology, surgery, and other.


¹⁰ Nyarko (2010) points out that a large fraction of the highly-skilled migrants in the United Kingdom from Southern and Eastern Africa are Asian or white according to the 2001 census.

¹¹ Naming data can be found at <http://www.census.gov/genealogy/www/data/2000surnames/index.html>

3. Estimation Methods

Our goal is to estimate location of birth, location of training, and age at migration for all doctors born and/or trained in Africa who were practicing in the United States at the end of 2010. As described above, the AMA Masterfile includes complete data on location of training for all registered doctors but only partial data on location of birth and no data on age at migration. The Venn diagram (Figure 1) helps to visualize what data are available from the AMA.

Figure 1: Venn diagram of the AMA dataset for location of birth and training



The left circle represents the doctors who were born in Africa and the right circle those trained in Africa. Their intersection, areas B and E, represents the doctors born *and* trained in Africa. The areas *above* the horizontal line denote the doctors who reported their country of birth in the AMA Masterfile while the areas *below* the line are those who did not answer that question. In other words, we know both the location of birth *and* training for those doctors in areas A, B, and C. On the other hand, doctors in areas E and F are in our file; we know that they were trained in an African country but we do not know their actual country of birth. This implies that we cannot be sure about the birth of doctors in areas E or F; hence the dashed line between them. Finally, area D represents doctors who were born in an African country but trained outside Africa *and* did not report their country of birth to the AMA. These people are completely missing from the AMA file and are indistinguishable from close to half a million other doctors in the database who did not report their country of birth.

The American Community Survey (ACS) provides complete data on location of birth and age at migration (along with numerous other demographic variables) but only for a nationally representative sample of doctors. Unfortunately, the ACS does not report country of education. To put it differently, we observe a

sample of the doctors in the circle on the left (areas A, B, D, and E) but we do not know if these people are in areas $A \cup D$ or they are in $B \cup E$ —i.e., trained outside or inside Africa.

Our key objective is to combine these different dimensions of information from the AMA Masterfile and the ACS survey to construct a more precise picture of high-skilled migration patterns. We need to identify the location of birth, location of training and age of migration for these doctors by matching the data as they are presented in the different datasets. To do so we proceed in three steps:

1. In the AMA data, we estimate the probability that an individual was born in each possible location of birth for those who do not respond to the country of birth question (areas E and F in Figure 1). We estimate these probabilities via propensity score matching relative to similar individuals who do respond (areas B and C).
2. We use the ACS data to estimate the size and distribution across countries of “missing doctors” in the AMA data (area D). We first use totals from the ACS data to generate an envelope that allows us to calculate the size of area D for each country of birth. Then we distribute these missing doctors across (non-African) locations of training by matching ACS observations to similar AMA observations with location of training data.
3. Finally, we estimate age-at-migration distributions. For African-trained doctors, we match the AMA data (including estimated country of birth probabilities from step 1) to the ACS data to estimate age at migration for each doctor in the AMA data. For doctors trained outside Africa, we simply use the ACS data with estimated location of training from step 2.

3.1. Trained in Africa

Consider a doctor i who lives in the United States, was born in location B_i and received medical education in location E_i . Define a series of indicator variables for whether that individual was born in location b and educated in location e .

$$D_{ibe} = I\{E_i = e \text{ and } B_i = b\}$$

For some individuals in the AMA data, D_{ibe} can be observed but we do not observe it for others due to missing location-of-birth data. As a result, the total number of immigrant doctors born in b and trained in e , N_{be} , includes an observed and unobserved component:

$$N_{be} = \sum_{i|D_{ibe} \text{ observed}} D_{ibe} + \sum_{i|D_{ibe} \text{ unobserved}} D_{ibe} = \sum_{i|D_{ibe} \text{ observed}} D_{ibe} + N_{be}^*$$

The latter term N_{be}^* cannot be calculated due to the missing data. So, we estimate N_{be}^* by matching AMA data on location of birth to observations missing such data. For each observation with missing country of birth, we estimate:

$$\hat{p}_{ib} = \Pr[B_i = b | \widehat{E}_i = e, X_i]$$

We employ standard kernel-based propensity score matching techniques (Heckman, Ichimura, and Todd 1998). It involves two steps. First, we estimate propensity scores—i.e., the probability of not reporting country of birth conditional on observable characteristics. We estimate this as a logit regression using the following covariates: gender, logarithm of age, an MD/DO dummy, a dummy for having an Asian name, a set of census division (regional) dummies, and a set of specialty group dummies. We stratify the matching by each location of training, and thus different coefficients are estimated for each country of training. Second, observations for which location of birth is not reported (regions E and F in Figure 1) are matched to observations for which location of birth is reported (regions B and C). This exercise is performed only for those doctors trained in Africa, and we match using an Epanechnikov kernel with bandwidth 0.06, again stratified by location of training. This process generates estimates of a probability distribution over country of birth for each observation with missing country of birth (i.e., \hat{p}_{ibe} estimates for each possible b for each observation with missing data). A small number of observations are dropped from the propensity score matching because a dummy variable leads to a unit or zero probability of reporting country of birth for those observations or to ensure common support. In the calculations we assume that their distribution of country-of-birth probabilities is at the average of others in their country of training.

Having generated these estimates, we can calculate the number of doctors born in b and trained in e as:

$$\hat{N}_{be} = \sum_{i|D_{ibe} \text{ observed}} D_{ibe} + N_{be}^* = \sum_{i|D_{ibe} \text{ observed}} D_{ibe} + \sum_{i|D_{ibe} \text{ unobserved}} \hat{p}_{ib}$$

This is the final result for doctors trained in location e . Repeating this process (for all 13 locations of education in Africa) yields estimates of the number of doctors born in any of the 19 locations and trained in one of the 13 African locations.

To generate standard errors for these estimates (and estimates described in future sections), we bootstrap the propensity score estimation procedure using 100 replications. Simple propensity score matching methods such as 1-to-1 matching cannot be bootstrapped because of the discontinuity in the matching process. However, kernel-based matching relies on continuous weights and thus does not suffer from this particular problem. As such, the literature has generally supported using bootstrapping for standard errors of kernel-based estimates (Abadie and Imbens 2008).

3.2. Trained Outside Africa

We use the estimation in section 3.1 to fully construct the place of birth for doctors who are trained in Africa (regions B, C, E, and F in Figure 1). However, there is still the issue of doctors who were born in an African country but trained outside of Africa and declined to report their country of birth to the AMA. As mentioned earlier, this group (area D in Figure 1) is not included in the extract from the AMA Masterfile used in this paper. We simply cannot observe their demographic and education information or even know how many such individuals exist because they are observationally identical to the larger population of doctors trained in the United States, Europe, etc. who also refused to report their country of birth. In order to identify this group, we combine the AMA Masterfile data with the 2010 file of the

ACS for all individuals who list “doctor” as their occupation and who report a birthplace in Africa.¹² While the absence of education location information and the small sample make this dataset potentially less reliable in counting African doctors, the data can be used to estimate how many African-born doctors are missing from the AMA data extract—i.e., the size of area D in Figure 1. We do this in three steps. First, we use ACS totals to provide an “envelope” for the number of immigrant doctors born in any particular country. This corresponds to areas A, B, D, and E in Figure 1. Since we have the estimates of areas A, B, and E from the AMA data and the previous section, we can determine the size of area D for each country of birth in Africa. Second, we match the ACS data to the AMA data to determine how those “missing doctors” should be distributed across countries of training outside Africa. Third, we implement a Bayesian correction to the estimates that allows us to improve the estimates using variables not present in both datasets.

Subject to sampling variation, the ACS data can measure the total number of doctors born in an African country who are in the United States. Combined with the estimates from the AMA data (see Section 3.1), we can estimate the number of doctors born in a given country but trained outside of it as:

$$\sum_{e|e \notin Africa} \hat{N}_{be} = \sum_{i|B_i=b} \omega_i - \sum_{e|e \in Africa} \hat{N}_{be}$$

The left side is the previously unknown number of doctors trained outside Africa but born in country b (i.e., areas A+D); ω_i is the ACS sample weight of observation i which is used in the first right-hand side term to estimate the total number of doctors born in country b . The final right-hand side term uses the estimates from the previous section to net out the number of doctors trained within Africa. So, we obtain an estimate for the total number of doctors born in any African country (areas A+D). An estimate of the number of missing doctors (area D) can be obtained by subtracting off the number of doctors in the AMA data who are trained outside Africa and report b as country of birth. Due to sampling variation, this value for “missing doctors” is slightly negative in two cases. For these two countries, we assume that there are no missing observations in the AMA data—i.e., we estimate the total number of doctors trained outside of Africa but born in these countries as the number of AMA respondents directly stating so.

This envelope method provides us with totals for the number of doctors born in each country and trained outside of Africa. Next, we distribute these totals among different locations of training. We perform a similar propensity score matching exercise to the one described above, but we use the ACS data and flip the roles of location of birth and location of education. Since the location of education is not observed for individuals in the ACS data, we match ACS observations to individuals in the AMA data who were born in the same location (conditional on reporting country of birth). In other words, the ACS

¹² There are some individuals for whom country of birth is vague. We categorize these with our “other” regional groups when possible. For those listed as “Africa, not elsewhere counted/not specified” we omit them from the propensity score matching but include them in the aggregate calculations by distributing them to the different countries of birth according to the empirical distribution for observations where the location of birth is clear.

observations now will be matched to areas A and B in Figure 1. All other details stay the same as above, and we obtain estimated probabilities across location of training for each ACS observation:

$$\hat{p}_{ie} = \Pr[E_i = e | \widehat{B}_i = b, X_i]$$

In the final stage, we implement a Bayesian correction to these propensity score estimates. Due to the limited overlap between variables in the ACS and AMA datasets, the propensity score estimates reflect matching on only three variables: age, gender, and census region of residence. While most of the available variables in the ACS must thus be discarded, it is possible to improve the estimated location of training probabilities by incorporating data on age of migration. In particular, we impose the restriction that

1. If an ACS observation reports that an individual migrated to the United States at age 18 or earlier, then it is the case that she was trained in the United States.
2. If an ACS observation reports that an individual migrated to the United States at age 30 or later, then it must be the case that he was trained outside the United States.

These corrections are supported by the data. In the AMA Masterfile, there are no doctors who graduated from medical school before the age of 18. Similarly, if an individual migrated to the United States at age 30 or older, and then went to medical school, he would graduate, at the earliest, at age 34. There are few African doctors in the AMA Masterfile who were born in Africa and graduated from an American medical school after age 34. So, we assume that doctors arriving at age 30 or older were educated outside the United States. While this information cannot be directly matched to the AMA data, we implement a Bayesian correction. We take the matching estimates \hat{p}_{ibe} and update these probabilities to account for the age at migration data. In particular, define \tilde{p}_{ibe} as the updated probability and A_i as the age at migration. Then, probability of being educated in the United States is:

$$\tilde{p}_{i,US} = \begin{cases} 1 & \text{if } A_i \leq 18 \\ 0 & \text{if } A_i \geq 30 \\ \hat{p}_{i,US} & \text{otherwise} \end{cases}$$

Conversely, the probability of being educated in country/location b outside the United States is then:

$$\tilde{p}_{ie} = \begin{cases} 0 & \text{if } A_i \leq 18 \\ \frac{\hat{p}_{ie}}{1 - \hat{p}_{i,US}} & \text{if } A_i \geq 30 \\ \hat{p}_{ie} & \text{otherwise} \end{cases}$$

where $\hat{p}_{i,US}$ is the original estimated probability that observation i was educated in the United States. Updating our estimates with the age at migration information helps improve the estimates and counteract the limited overlap between AMA and ACS variables.

Finally, we use the totals from the envelope and the updated probabilities \tilde{p}_{ie} to generate estimates of the aggregate number of doctors born in an African country b and trained in a non-African country e . We use the ACS data to calculate the aggregate probability of being educated in location e (given that location of birth is b) as:

$$\tilde{p}_{e|b} = \frac{1}{N_b} \sum_{i|B_i=b} \tilde{p}_{ie} * \omega_i$$

where ω_i is the ACS survey weight of observation i and N_b is the sum of survey weights of ACS observations born in location b . We then use these probabilities to distribute the envelope totals across potential locations of training. For a non-African location of training e :

$$\hat{N}_{be} = \left(\sum_{i|B_i=b} \omega_i - \sum_{k|k \in Africa} \hat{N}_{bk} \right) * \frac{\tilde{p}_{e|b}}{\sum_{k \notin Africa} \tilde{p}_{k|b}}$$

These results provide estimates for the number of doctors born in an African location and trained at a non-African location for all such locations—i.e., country-by-country estimates of area A+D in Figure 1. This completes joint estimation of country of birth and country of education for African immigrant doctors.

3.3. Estimating Age at Migration

The previous two sections describe how we estimate the relationship between location of birth and location of training. Adding estimates of when African doctors migrate to the United States would provide another critical piece of information about their career paths. The main complication is that the ACS data has information on age at migration and location of birth, but it lacks information on location of training, as noted previously. Meanwhile, the AMA data lacks data on age at migration though it has the location of training. We combine these data again to estimate the age of migration distribution for each location of birth location and training (b,e) pair. There are three groups of doctors we need to be concerned about and we approach each group differently:

1. For doctors born and trained in Africa (areas B and E in Figure 1), we estimate age at migration in the AMA data by matching it to ACS observations.
2. For doctors born in Africa but trained outside (areas A and D), we simply use the ACS data and the matched location of training from the previous section. We use separate methods for groups 1 and 2 to ensure that our age at migration estimates are based on the same data as the birth and training location estimates.
3. For doctors trained in Africa but born outside Africa (areas C and F) we cannot estimate age at migration. These doctors only appear in the AMA data but cannot be observed in the ACS and thus their age at migration cannot be estimated.

3.3.1. Estimating Age at Migration for Those Born and Trained in Africa

Since the AMA Masterfile does not include any information on age of migration, we must estimate it by matching the AMA observations to similar ACS observations. As before, the matching occurs only for observations within the same country of birth. In particular, the AMA data includes two groups of people: those who report country of birth and those who do not. For those who did not report, we estimated a probability distribution for country of birth over the 19 geographic locations in the first section. When matching in this section, we include all AMA observations *for which this probability is greater than zero* and match them to ACS observations born in country b . As before, we perform a kernel-weighted matching based on gender, log of age, and census division dummies, and we estimate the probability that age of migration for individual i :

$$\hat{m}_{ai} = \Pr[A_i = a | X_i, \widehat{\Pr}\{B_i = b\} > 0]$$

This probability can be calculated for any value of a to estimate the entire probability distribution of age at migration for i .

As with Section 3.2, matching between the ACS and AMA data can be improved by incorporating information beyond those variables that directly correspond between the two datasets. In particular, the AMA data contain complete information on location of training and age at graduation from medical school, and this information can be used to apply a Bayesian correction similar to the one above. More specifically, we assume that:

1. Receiving training in the United States implies that age at migration is less than or equal to age at graduation from medical school.
2. Receiving training outside the United States implies age at migration is greater than or equal to age at graduation from medical school.

Formally, define G_i to be age at graduation. Thus, if individual i is trained in the United States, we update the probabilities:

$$\tilde{m}_{ai} = \begin{cases} 0 & \text{if } a > G_i \\ \frac{\hat{m}_{ai}}{1 - \sum_{a > G_i} \hat{m}_{ai}} & \text{if } a \leq G_i \end{cases}$$

If observation i is trained outside the United States then we apply the reverse:

$$\tilde{m}_{ai} = \begin{cases} 0 & \text{if } a < G_i \\ \frac{\hat{m}_{ai}}{1 - \sum_{a < G_i} \hat{m}_{ai}} & \text{if } a \geq G_i \end{cases}$$

Together, this procedure updates each individual's matched age at migration distribution to reflect information about age at graduation. A very small number of observations are dropped due to a zero

denominator (i.e., there are no observations from their country of birth in the ACS data with age at migration consistent with their age of graduation). As before, we aggregate these probabilities across individuals to estimate the age at migration distribution for any given pair of country of birth and country of education.¹³ A similar process can be applied to a particular cohort by simply summing over those in the appropriate age range.

3.3.2. Estimating Age at Migration for Those Trained Outside Africa

Estimating age-at-migration distributions for those trained outside Africa requires using the ACS data but requires no new analysis beyond Section 3.2. Location of birth is observed in the ACS data and probabilities for different locations of training were estimated in Section 3.2. Age at migration, A_i , is also recorded in the data for all individuals. Thus, we can estimate the number of people that were born in location b , educated in location e , and migrated to the United States at age a by simply counting up observations with the appropriate age at migration and country of birth, weighting them by the survey weights and the probability that the individual was trained in location e . This count can then be turned into a probability by dividing by the total number of doctors born in b and trained in e .

3.3.3. Combining Age at Migration Distributions

We then combine the results from Sections 3.3.1 and 3.3.2. To be consistent with our previous estimates, we use the ACS-based estimates from 3.3.2 to generate age at migration distributions when the country of training is outside Africa. Meanwhile, we use the AMA-based estimates from 3.3.1 for those with training inside Africa. As noted above, these estimates are only for those born in Africa. Unfortunately, we cannot estimate age at migration distributions for those born outside of Africa but trained in Africa. Finally, with these estimates in hand, we can convert the probabilities into age at migration cumulative distribution functions for each location of birth-location of training pair.

4. Results

The previous section describes the three stages of the empirical strategy to construct a more precise profile of migrant African doctors in the United States in terms of their location of birth, location of training, and age of migration. This section presents these results in the same sequence.

4.1. Propensity Score Matching Results

The first stage involves estimating the location of birth probabilities for the 19 regions for those who did not report this information in the AMA data (areas E and F in Figure 1) by matching them to those trained in the same region who reported this information (areas B and C). Table 3 presents the main coefficients from estimating the propensity score by country of training. The displayed marginal effects

¹³ This time, though, it is necessary to weight each observation by the estimated probability that the observation was in fact born in location of birth b .

indicate how the probability of not reporting country of birth changes with observed characteristics. The estimated coefficients are then used to compute propensity scores and complete the matching process, as described above. For brevity, we do not report the coefficients of the medical specialization, census region and DO dummy variables.

The second stage of our estimation involves the “missing doctors” from the AMA dataset (area D in Figure 1) who were born in Africa, trained elsewhere and did not report their country of birth to the AMA. In order to estimate the size and location of birth of this group, we use the ACS data as envelope to determine the size of area D and match the ACS and AMA data as described in the previous section. The results of the propensity score estimation are reported in Table 4. Again for space reasons, the marginal effects for census regions in the United States are not reported. We should note that gender coefficients are not significant in most of the cases, indicating that the migration and education patterns of male and female physicians are quite similar. This is an important distinction with low-skilled migration patterns and deserves additional scrutiny.

Our third and final estimation exercise determines age at migration by matching the AMA and the ACS data one last time. We can perform this estimation only for those born in Africa since migration age comes from ACS and we cannot identify those born outside but trained in Africa in the ACS. Recall that, in the AMA data, we have a probability distribution over the birth location for every individual after the first stage (see above). This is basically a vector of length 19. If the individual reported his birthplace, one of these entries is equal to one and the rest are zeros. If he did not report and we estimated probabilities, these 19 entries add up to 1. For every one of the 13 African birth locations, we take the individuals with non-zero probability of being born there and match them to the observations with the same birth country in the ACS data. The results of the estimation of the propensity scores from this process are reported in Table 5. The marginal effects of census regions are not reported.

After the matching, we perform adjustments as discussed above and weigh matched age at migration dummies using both the ACS population weights and the probability weights from our first stage estimation. Our end-result is a prediction over the age of migration for each individual. We then sum up these probabilities to determine what percentage of the doctors born in location i and trained in location j migrated to the United States at a given age.

4.2. Missing Location of Birth and Education in the Databases

This section presents the place of birth and education matrices from the first stage of our estimation exercise. We start with those who actually report their location of birth to the AMA. Table 6 presents the matrix of location of training (columns) against location of birth (rows) for those 8,062 doctors who reported their country of birth. For example, we see that a total of 2,520 people reported Egypt as their birthplace and 1,816 of these were actually trained in Egypt while 558 were trained in the United States.

Table 6 does not include the 47 percent of the AMA sample (7,129 doctors) who did not report their country of birth; our predictions on their location of birth are reported in Table 7. Continuing with the Egyptian example from above, we note that there were 1,964 Egyptian-trained doctors who did not report birth information and our estimation predicts that 1,712 of these were born in Egypt. An

additional 252 of them were estimated to be born outside Egypt, mostly allocated to the “rest of the world” category that includes Arabic-speaking countries in the Middle East. We should note that the last six columns (non-African regions) are left blank in this matrix as it would include our predictions for people who were born in Africa but trained in the rest of the world. As mentioned earlier, these are not included in our AMA sample unless country of birth was reported. Finally, Table 8 combines Tables 6 and 7 to give a full picture of the AMA data after our first stage predictions are completed.

Table 8 reveals our first set of important results. Of the doctors recorded in the AMA database, only 81 percent of those trained in an African country were born in the same country. Around 4 percent were born in other African countries and a staggering 15 percent (1,730 doctors) were born outside Africa but trained there. This number includes 539 doctors who were born in the United States, went to medical school in Africa and then returned to the United States to practice their profession.

The second stage of our estimation predicts the number of doctors who were born in Africa, trained outside Africa and did not report their country of birth to the AMA. Table 9 reports the resulting matrix of allocating these missing doctors from 13 birth regions in Africa to the 6 non-African regions of training. We predict that almost 70 percent (6,000 out of 8,720) of these doctors were trained in the United States, most likely children who migrated with their parents and students who came to attend medical school.


Our final step is to combine these two stages in order to construct an overall matrix of training and birth. To this end we use estimates from the AMA data when the location of training is one of the 13 African regions (the first 13 columns of Table 8) and from the ACS predictions for the six non-African regions of training (the last six columns of Table 9). The resulting final matrix, our ultimate objective, is presented in Table 10.

4.3. Patterns in Location of Birth and Education

We identify several key patterns that arise these results, especially in Table 10, which should provide insights on global skilled-migration flows. The education and birth matrices are especially useful in identifying where the largest differences are likely to exist between the numbers for locations of birth and of education.

First, in both the aggregate and, for many individual countries, the measured magnitude of migration numbers vary widely depending on whether country of birth or country of training is used as the main criteria. Our empirical exercise estimates that there are 19,997 African-born and/or -trained doctors in the United States (Table 10). Of this group, 9,547 (47.7 percent) were born *and* trained in Africa. On the other hand, 8,720 (43.6 percent) were born in but trained outside Africa and 1,730 (8.7 percent) were born outside but were trained in Africa (see Figure 2). In other words, there are 11,277 doctors trained and 18,267 doctors born in Africa. The overall size of medical migration from Africa to the United States would be more than 60 percent greater if we were to use place of birth instead of education criteria. However, the main value of our exercise arises from the joint prediction of the birth and education dimensions in this discussion.

Figure 2: Distribution of African-born and -trained doctors in the United States



In Table 11, we present the details of the information from Table 10 in a compact format for the five African countries with the largest number of doctors who were trained there: Egypt, Ethiopia, Ghana, Nigeria, and South Africa. Jointly, doctors trained in these five countries account for almost 65 percent of the entire AMA sample used and 87 percent of migrant doctors trained in Africa. Our empirical exercise reveals further interesting patterns, especially the stark differences between countries on the same continent. Of the 4,062 Egyptian-trained doctors, the largest group (3,528) was born in Egypt (87 percent). We predict that 365 doctors born outside of either Africa or the United States, mostly in other Arab countries, were also trained in Egypt. The two smallest countries of this group are Ghana and Ethiopia, and they train very few foreign-born doctors; almost all of the doctors trained in these countries were also born there. On the other hand, almost 40 percent of the doctors trained in South Africa and over 20 percent of those trained in Nigeria are born outside their borders. Both of these countries are important training hubs for other African medical students, as well as for many students from the United States and other parts of the world.

Thus, our second result shows that a non-negligible number of doctors are born outside Africa, train in Africa, and then migrate to the United States to work; a pattern that has not been identified before. There are 1,730 such doctors, comprising slightly over 15 percent of the 11,277 doctors trained in Africa. Almost one-third of this group (532 people out of 1,730) were born in the United States; 14 percent (237 people) were born in other English speaking countries, 13 percent (225 people) were born in South Asia, and 11 percent (182 people) were from Europe. These patterns indicate that the mobility of highly-educated professionals starts at a very young age and we need careful distinction between birth and training locations to identify such dynamics. Most analyses start with the assumption that African skilled migration is a one-way flow. Our results, however, show that African medical schools attract a non-

negligible number of outsiders despite their limited resources. This fact has significant implications for health-related education and human resource policies in Africa.¹⁴

Our third result focuses on the location of education of the African-born doctors. Of the 9,547 doctors born and trained in Africa, 9,111 (over 95 percent) were trained in their birth country and slightly over 400 moved internally within Africa for their education. When we look at those born in but trained outside of Africa, we see that a majority of these 8,720 doctors chose the United States for their education (almost 70 percent), as shown in Table 10. The age at which these people arrived in the United States is explored further in the next section. At the outset, we would expect many African doctors to go to European schools due to colonial linkages and relative geographic proximity. Around 12 percent chose Europe and other non-US English speaking OECD countries for their medical schools. Surprisingly, large numbers (8.5 percent) go to South Asian countries. Such students tend to be descendants of the South Asian migrants who went to East and South Africa during the British colonial rule of the previous century. These numbers prove that diaspora bonds remain relatively strong, as the grandchildren of Indian migrants of the 19th and early 20th centuries go back to their grandparents' home countries to study. Like their grandparents, they choose to cross an ocean to settle in a distant land.

In Table 12, we look at the largest five countries in terms of the numbers of doctors born and we clearly see the importance of predicting the “missing” doctors from the AMA database. These are the same countries as before: Egypt, Nigeria, South Africa, Ghana, and Ethiopia in descending order of numbers. According to our estimates, there are 4,332 doctors in the United States who were born in Egypt, and as noted earlier, 3,528 (over 81 percent) were trained at home. However, there are over 800 Egyptian-born doctors trained outside Egypt, most of them trained in the United States. Similar statistics are significantly larger for the sub-Saharan countries. For example, fewer than half of the doctors born in Ethiopia or South Africa and practicing in the United States were actually trained in their respective birth-countries—just 44 percent of the Ethiopian-born and 47 percent of the South African-born. Furthermore, 17 percent of Ethiopian-born, 14 percent of South African-born, 13 percent of Nigerian-born, and 11 percent of Ghanaian-born doctors were trained outside of Africa and the United States. This “transit” migration is an additional dimension of global skill flows that deserves more attention than it has received to date.


Our last notable observation is that around 7 percent of the group that trained outside Africa went to the Caribbean countries (see Table 10). Since medical school capacity has not kept up with the increased demand for doctors in the United States, new private medical schools emerged in various Caribbean countries. There is a significant debate on the quality and value of these medical schools as they tend to attract students who seem to have failed to get into medical schools in the United States. Our data indicate that American students are not the only clients of the Caribbean medical schools; significant numbers of African-born doctors are enrolling there, probably with the intention of eventually migrating to the United States. There is a large “brain gain” literature, starting with Stark, Helmenstein, and

¹⁴ It is important to know whether these students born outside Africa pay higher fees. In some countries they do but there is no comprehensive data or analysis on this issue that we know of.

Prskawetz (1997) and Mountford (1997), on the incentives to acquire human capital through increased migration opportunities. African born doctors—mostly from Nigeria and other West African countries—who go to schools in the Caribbean and practice in the United States provide quite interesting evidence for this argument. Of course this cannot be truly considered brain gain for Africa, as defined in the literature, since these students do not return home but migrate to another destination.

Our final discussion in this section will use the case of doctors from South Africa to illustrate the importance of the distinctions discussed above. Who really is a South African doctor? Figure 3 provides a nuanced answer to this difficult question using the numbers from Tables 11 and 12 for South Africa. We see that there are 818 doctors born and trained in South Africa who later moved to the United States, but this represents only a minority of the 2,642 doctors who either were born or trained there. By contrast, there are 1,002 doctors born in South Africa but trained in the United States, and another 290 trained in other non-African countries around the world. Meanwhile, South Africa trains a significant number of doctors from other African countries (181) and the United States (92) as well as other countries in the world (254). Our analysis enables us to split the numbers further. We can identify the other eight African regions in which these 181 doctors were born, as well as the non-US and non-African countries from which the other 254 doctors originated.

Figure 3: Distribution of South African-born and -trained doctors



The standard errors of our estimates are provided below each number in Tables 11 and 12 for the largest five countries and Table 13 for the whole sample. As can be seen, all of the estimates are statistically different from zero except for a few cells with very low values—such as trained in Ethiopia and born in the rest of the world. More importantly, these small standard errors imply tight confidence intervals around estimates for the largest country-of-birth or country-of-training career paths for African immigrant doctors. For instance, the confidence interval for the number of doctors trained in Egypt but


born outside Africa or the United States is (340, 390). Thus, our data provide precise estimates that allow us to examine the interplay of birth and training locations in a meaningful way. But does this vary across countries? In particular, the Nigerian column shows the highest standard errors due to the fact that Nigerian-trained doctors are much less likely to report their country of birth. While there might be many cultural and political reasons for this reluctance, the effect is to cause larger standard errors in our results.

4.4. Age of migration

The time spent at home after graduation but before migration to the United States is an important dimension in the career path of a migrant doctor. It also has important welfare and policy implications for both the origin and destination countries. If migration takes place at a relatively young age, it means most of the benefits of the accumulated human capital will be enjoyed by the consumers in the destination country and “lost” to the country of training. On the other hand, if the migrant doctors practice their profession for some time before migrating, then the benefits from human capital accumulation will be more evenly split between countries of education and destination.

Naturally, the age of migration needs to be predicted separately for each doctor but we report the results separately for each location of training and birth pair. It would not make sense to combine Egyptian-born doctors who are trained at home with those trained in the United States, since the latter group is more likely to have arrived as children and students. We start with Figure 4 which presents the cumulative distribution functions of all African-born doctors in the United States. The blue line represents the distribution of age-at-migration to the United States for all doctors, regardless of their country of training. The other lines split this out by location-of-training at home (red), in the United States (green), or somewhere else (purple). First, we note that the age of migration profile is quite different for these three groups, as expected. These differences once again stress the importance of differentiating high-skilled migrants based on where their human capital was obtained. Among those trained in the country of birth, the median age at migration is 30 years, and 90 percent arrive by age 39. However, those trained in the United States arrive much earlier; the median age of migration is only 23 years, and 90 percent arrive by age 29. About 30 percent arrive as children and most of the remainder arrived during the 18-29-years age range, since their migration is likely to be motivated by schooling. Finally, those training elsewhere appear similar to those trained in their birth countries.


Figure 4: Age of migration, by location of training
(cumulative probability distribution)




The next step is to compare the migration behavior of the doctors trained in the main five countries we analyzed earlier: Egypt, Ethiopia, Ghana, Nigeria, and South Africa. Figure 5 presents results for those doctors trained at home. Among those trained in Egypt, the median age of migration to the United States is 29.5 years, over 80 percent of them arrive by the age of 34, and 90 percent arrive by age 38. Doctors born and trained in South Africa exhibit almost the same patterns as Egyptian doctors. The median age of migration is around 30 years, and 90 percent migrate by the age of 38. Nigerian-born doctors show a slightly different pattern; for those trained at home, the median age of migration is 32 years, and 90 percent of migration is completed only by age 40. In other words, Nigerian doctors who were trained at home delay their migration significantly compared to the other large groups. Our final group is composed of doctors born and trained in Ghana, who seem to migrate to the United States relatively sooner after graduation. Their median age of migration is 28 years, and 90 percent migrate by the age 32.

Our next comparison is between doctors trained in the United States but born in the five major countries. Figure 6 shows that Egyptian-born doctors migrate earliest, at a median age of 17 years. Nigerian doctors are the slowest to migrate, at the median migration age of 24 years. This pattern indicates that most of the Nigerian doctors migrate specifically to attend medical school, while Egyptian doctors are more likely to migrate as children with their families. Among the other three countries, the migration patterns are more gradual, indicating a mixture of these two processes.

**Figure 5: Age of migration of doctors trained in one of five countries
(cumulative probability distribution)**




**Figure 6: Age of migration of doctors trained in the US but born in one of five countries
(cumulative probability distribution)**



The final graph presents the age-at-migration analysis for different age cohorts of doctors born and trained in sub-Saharan Africa (Figure 7). There are three age cohorts analyzed by their current age: 35-45 years, 45-55 years, and 55-65 years. The critical observation is that younger (more recent birth) cohorts seem to be arriving at an earlier age; the distribution functions are shifting to the left. For example, the median age at migration is declining by one year with each cohort, from 31.5 years to 29.5 years. Given that the average service at home is only 7 years (assuming a graduation age of 22 years), this is a considerable decline. By comparison, we do not observe the same shift among the doctors from Egypt and other North African countries.

Figure 7: Age of migration of doctors from and trained in Sub-Saharan Africa, years (cumulative probability distribution)



Different age-of-migration profiles have strong implications for education and labor market policies. For a country such as Ghana, the classic story of medical brain drain may be accurate, at least for those doctors who train at home. Ghanaian doctors tend to leave quickly after completion of medical school, giving the public health system little time to recoup benefits from the public subsidy of education. On the other hand, doctors from Nigeria and South Africa tend to wait longer between completing their education and migrating to the United States. Although this is not observed in our data, it is likely that this corresponds to time spent working in the country of training. As a result, public health systems in these countries may well benefit from several years of local medical service by the doctor prior to migration.

5. Conclusion

There are over 20 developing countries where more than half of their tertiary-educated citizens are abroad (Artuc et al. 2015). Such widely-cited high-skilled migration or “brain drain” numbers are inflated by our reliance on census-type data sources where migrants are classified according to their countries of birth or citizenship, rather than their countries of education. If education was completed in the destination (or a third country) and/or if the emigration took place at a later point in a professional’s career, then the welfare and policy implications for both the origin and destination countries would be quite different.

We explore these nuances by focusing on the African-born and/or -trained physicians who are currently practicing in the United States. We use propensity score matching techniques to merge education data from AMA data with country-of-birth and age-of-migration data from the ACS. In the end, we predict the location of birth, location of education, and age of migration for close to 20,000 doctors. We find quite surprising patterns.

First, around 48 percent of these doctors were born and educated in Africa, mostly in their home countries. Second, another 44 percent were born in Africa but were educated outside the continent, including the Caribbean to South Asia. Finally, African medical schools trained 1,730 doctors who were born outside Africa. In addition to identifying location of birth and education patterns, we determine the age of migration profiles of various migrants and highlight widely heterogeneous patterns across birth and training countries. How long various professionals, especially doctors, stay at home and work before they migrate to the other countries is an important human resource issue for policymakers. We perform the same analysis by age cohorts and find that more recent cohorts have migrated at an earlier age, indicating the time spent at home after graduation is shrinking.

Overall, our results indicate that skilled migration varies widely across countries. A country such as Egypt sends many native-born and -trained doctors abroad but also trains students from other countries. In such a context, policymakers may reasonably discuss whether admitting foreign students might offset some of the financial loss incurred when native students receiving education subsidies migrate. A country such as Ethiopia faces a different situation: only two in five Ethiopian-born doctors in the United States were trained in Ethiopia. Policymakers in this situation should avoid the factual error of conflating Ethiopian-born doctors in the United States with Ethiopian-trained doctors and subsequent lost educational subsidies. Nonetheless, such a fact suggests that public policy aimed at recruiting more doctors in Ethiopia may reasonably need to focus on why the families of talented children emigrate.

There are three other important issues in the high-skilled migration literature that this paper cannot address. The first concerns financing of education. While medical education is publicly financed in most countries, it is generally privately-funded in the United States, and especially so in the Caribbean. However, there are many exceptions. Some schools in Africa charge tuition to cover their costs, especially for foreign students. Many African countries also provide scholarships to their students to study abroad, especially in the developed world, with the expectation that they will return home (Clemens and Petterson 2008). The second issue is retention of high-skilled workers. Antwi and Phillips

(2013) show that higher wages can reduce emigration of doctors trained in Ghana. But we need detailed information on doctors' and students' demographic and personal characteristics to assess the impact of various policy interventions. Third is the issue of brain waste. Many people who were trained as doctors in their home countries fail to obtain a medical license or choose not to practice in the United States. As a result, they do not register with the AMA or appear in the ACS sample. The extent and causes of brain waste have important policy implications (Mattoo, Neagu, and Özden 2008). Unfortunately, existing administrative and survey-type data are not sufficient to address these funding, retention, and brain-waste questions. Ideally, future work should combine the results and methods of the present paper with more detailed data from personal surveys of doctors to fully answer these questions.

References

- Abadie, A. and G.W. Imbens. 2008. "On the Failure of the Bootstrap for Matching Estimators." *Econometrica*, 76(6): 1537-1557.
- Antwi, J. and D.C. Phillips. "Wages and Health Worker Retention: Evidence from Public Sector Wage Reforms in Ghana." *Journal of Development Economics*, 102 (2013): 101-115.
- Artuc, E., F. Docquier, C. Özden, and C. Parsons. "A Global Assessment of Human Capital Mobility: the Role of non-OECD Destinations, *World Development*, 65 (2015): 6-26.
- Beine, M., F. Docquier, and H. Rapoport. 2007. "Measuring International Skilled Migration: New Estimates Controlling for Age of Entry." *World Bank Economic Review*, vol. 21: 249-254.
- Bhagwati, J. and K. Hamada. 1974. "The Brain Drain, International Integration of Markets for Professionals and Unemployment: a Theoretical Analysis." *Journal of Development Economics*, 1(1): 19-42.
- Bhargava, A. and F. Docquier. 2008. "HIV Pandemic, Medical Brain Drain, and Economic Development in sub-Saharan Africa." *The World Bank Economic Review*, 22(2): 345-366.
- Bhargava, A., F. Docquier, and Y. Moullan. 2011. "Modeling the Effects of Physician Emigration on Human Development." *Economics & Human Biology*, 9(2): 172-183.
- Chen, L. C. and J.I. Boufford. 2005. "Fatal Flows—Doctors on the Move." *New England Journal of Medicine*, 353(17): 1850-1852.
- Clemens, M. 2007. "Do Visas Kill? Health Effects of African Health Professional Emigration." *Center for Global Development Working Paper No. 114*.
- Clemens, M. and G. Pettersson (2008) "New data on African health professionals abroad." *Human Resources for Health*, 6(1).
- Docquier, F. and A. Marfouk. 2006. "International Migration by Educational Attainment 1990-2000." *International Migration, Remittances and Development*. Özden, C. and M. Schiff (eds.), Palgrave Macmillan, New York.
- Docquier, F. and H. Rapoport. 2012. "Globalization, Brain Drain and Development." *Journal of Economic Literature*, vol. 50, no.3: 681-730.
- Easterly, W. and Y. Nyarko. 2009. "Is the Brain Drain Good for Africa?" *Skilled Migration Today: Prospect, Problems and Policies*. Bhagwati, Jagdish and Gordon Hanson (eds.): 316–60. Oxford University Press.

- Gibson, J. and D. McKenzie. 2011. "Eight Questions about Brain Drain." *Journal of Economic Perspectives*. Vol.25. no.3: 107-28
- Hanson, G. 2010. "International Migration and the Developing World." *Handbook of Development Economics*, Vol. 5. Rodrik, Dani and Mark Rosenzweig (eds.). North Holland, Amsterdam: 4363-4414.
- Kangasniemi, M., L.A. Winters, and S. Commander. 2007. "Is the Medical Brain Drain Beneficial? Evidence from Overseas Doctors in the UK." *Social Science and Medicine*, 65(5).
- Lee J. 2006. "Message from the Director-General." *Working Together for Health: The World Health Report, 2006*. World Health Organization.
- Mattoo, A., C. Neagu, and C. Özden. 2008. "Brain Waste? Educated Immigrants in the US Labor Market," *Journal of Development Economics*, Vol.87, No. 2: 255-69.
- Mountford, A. 1997. "Can a Brain Drain Be Good for Growth in the Source Economy?" *Journal of Development Economics*, 53(2).
- Mullan, F. 2005. "The Metrics of the Physician Brain Drain." *New England Journal of Medicine*, 353(17): 1810-1818.
- Nyarko, Y. 2010. "EU Policies and African Human Capital Development." EUI-RSCAS Working Paper No.30, Robert Schuman Center for Advanced Studies, EUI, Florence.
- Okeke, E. 2009. "An Empirical Investigation of Why Doctors Migrate and Women Fail to Go For Screening." Unpublished dissertation, University of Michigan.
- Özden, C., C. Parsons, M. Schiff, and T. Walmsley. 2011. "Where on Earth is Everybody? The Evolution of Global Bilateral Migration 1960–2000?" *World Bank Economic Review*, 25(1): 12-56.
- Stark, O., C. Helmenstein, and A. Prskawetz. 1997. "A Brain Gain with a Brain Drain." *Economics letters*, 55(2): 227-234.

Table 1: American Medical Association (AMA) data on country of training and birth

COUNTRY	Fraction Trained	Fraction Born	Mean Age	Fraction Female	Mean Age at Migration	Fraction Asian Name
Total	1.00	1.00	47.0	0.27	26.0	0.05
Egypt	0.27	0.17	49.5	0.20	24.2	0.01
Ethiopia	0.03	0.04	44.7	0.17	25.1	0.01
Ghana	0.05	0.05	47.0	0.25	27.1	0.00
Kenya	0.01	0.05	46.4	0.22	25.5	0.22
Liberia	0.00	0.01	54.1	0.29	28.7	0.02
Nigeria	0.21	0.06	45.3	0.32	24.8	0.02
South Africa	0.09	0.03	52.8	0.19	24.1	0.02
Sudan	0.02	0.01	42.9	0.31	25.1	0.01
Uganda	0.01	0.01	44.9	0.42	26.3	0.14
Zimbabwe	0.01	0.01	46.2	0.25	24.8	0.08
Other East Africa	0.01	0.02	50.2	0.29	25.9	0.34
Other North Africa	0.03	0.02	42.7	0.32	26.0	0.01
Other West, Central, & Southern Africa	0.01	0.02	45.7	0.29	27.3	0.00
Caribbean	0.02	0.00	43.0	0.33	30.5	0.11
English Speaking	0.02	0.00	44.7	0.31	27.7	0.32
Europe	0.02	0.00	50.5	0.24	27.6	0.02
South Asia	0.02	0.01	54.4	0.28	25.0	0.69
United States	0.17	0.01	43.5	0.37	28.7	0.08
Rest of World	0.01	0.02	47.8	0.21	26.7	0.06
Missing	0.00	0.47				

Source: American Medical Association Physician Masterfile

Table 2: American Community Survey (ACS) data on country of birth & age of migration

COUNTRY	Fraction Trained	Fraction Born	Mean Age	Fraction Female	Mean Age at Migration
Egypt	--	0.24	47.8	0.19	27.5
Ethiopia	--	0.07	41.9	0.27	26.3
Ghana	--	0.05	42.8	0.22	24.5
Kenya	--	0.05	45.6	0.25	23.2
Liberia	--	0.02	47.4	0.30	25.6
Nigeria	--	0.24	43.0	0.35	27.5
South Africa	--	0.12	50.8	0.25	27.9
Sudan	--	0.02	44.2	0.09	29.8
Uganda	--	0.02	54.8	0.20	26.1
Zimbabwe	--	0.02	48.6	0.23	23.7
	--				
Other East Africa	--	0.06	44.2	0.35	23.0
Other North Africa	--	0.04	41.6	0.37	25.4
Other West, Central, & Southern Africa	--	0.05	38.9	0.24	25.5
Missing	--	--	--	--	--

Source: American Community Survey

Table 3: Propensity Score Coefficients—within AMA match for missing location of birth

Covariate	Location of Training												
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE
Female	0.015 (0.021)	-0.079 (0.067)	0.10** (0.048)	0.069 (0.11)	-0.19 (0.27)	-0.020 (0.016)	0.057 (0.12)	0.100* (0.057)	0.31** (0.15)	-0.0017 (0.031)	0.084 (0.064)	0.017 (0.10)	0.19 (0.16)
Log(Age)	-0.15*** (0.042)	0.43*** (0.15)	0.16 (0.12)	0.29 (0.28)	-2.16* (1.21)	-0.91*** (0.046)	-0.53 (0.34)	0.21 (0.15)	0.68* (0.41)	0.067 (0.069)	0.24 (0.17)	-0.70*** (0.24)	-0.94** (0.38)
Asian Name Dummy	-0.072 (0.10)	0.19 (0.19)	-0.20 (0.27)	-0.0015 (0.12)	--	-0.14** (0.062)	-0.0053 (0.12)	-0.15 (0.26)	--	-0.053 (0.084)	--	0.15 (0.12)	0.14 (0.20)
Census Division Dummies	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES
Specialty Dummies	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES
DO Dummy	YES	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Observations	4061	518	695	164	46	3174	116	430	78	1345	318	141	86
Pseudo R ²	0.01	0.08	0.03	0.12	0.25	0.14	0.17	0.06	0.15	0.03	0.06	0.11	0.33

The table reports average marginal effects from a logit model where the dependent variable is an indicator of whether data on the individual's location of birth is missing. Each column corresponds to a different sample, in particular the sample of individuals in the AMA data with a location of training in the listed country. The full range of covariates includes 10 census divisions and twenty specialty groups. Statistical significance at the 1, 5, and 10 percent levels are denoted respectively by *, **, and ***. Standard errors are in parentheses.

Table 4: Propensity Score Coefficients—ACS to AMA match for missing location of training

Covariate	Location of Birth												
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE
Female	-0.0098 (0.011)	-0.024 (0.022)	-0.011 (0.014)	-0.00078 (0.020)	-0.053 (0.062)	-0.0043 (0.021)	0.014 (0.029)	0.023 (0.033)	-0.023 (0.027)	-0.010 (0.042)	-0.087** (0.043)	-0.041 (0.040)	-0.0096 (0.087)
Log (Age)	-0.052** (0.020)	-0.060 (0.056)	-0.027 (0.032)	-0.018 (0.037)	0.036 (0.13)	-0.75*** (0.063)	-0.024 (0.052)	-0.097 (0.063)	-0.054 (0.059)	-0.016 (0.11)	0.025 (0.10)	0.30*** (0.11)	-0.39 (0.28)
Asian Name Dummy	--	--	--	--	--	--	--	--	--	--	--	--	--
Census Division Dummies	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES
Specialty Dummies	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
DO Dummy	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Observations	2713	600	852	710	106	1163	416	363	357	513	122	169	99
Pseudo R ²	0.01	0.02	0.03	0.01	0.02	0.2	0.04	0.03	0.07	0.01	0.06	0.09	0.07

The table reports average marginal effects from a logit model where the dependent variable is an indicator of whether the observation is in the ACS dataset (i.e. has missing location of training) . Each column corresponds to different samples defined by location of birth. Each column includes all ACS observations from the listed country of birth and all AMA observations that report the listed country of birth (those reporting no country of birth are omitted from all columns). The full range of covariates includes 10 census divisions. Statistical significance at the 1, 5, and 10 percent levels are denoted respectively by *, **, and ***. Standard errors are in parentheses.

Table 5: Propensity Score Coefficients—AMA to ACS match for estimating age at migration

Covariate	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE
Female	0.0043	0.0014	0.0057	-0.0044	-0.0046	-0.0097*	-0.0029	-0.0075	-0.00063	-0.0021	0.0052	0.016	-0.0038
	-0.0059	-0.0055	-0.0045	-0.0061	-0.007	-0.0059	-0.0052	-0.0049	-0.0016	-0.012	-0.0041	-0.019	-0.0088
Log (Age)	0.036***	0.053***	0.039***	0.034***	0.016	0.19***	0.039***	0.027***	0.020***	0.11***	0.012	-0.20***	0.022
	-0.012	-0.012	-0.012	-0.0098	-0.013	-0.015	-0.0099	-0.0081	-0.0046	-0.025	-0.01	-0.051	-0.016
Asian Name Dummy	--	--	--	--	--	--	--	--	--	--	--	--	--
Census Division Dummies	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES	YES
Specialty Dummies	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
DO Dummy	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Observations	5,102	2,175	2,248	2,498	1,081	3,947	2,065	2,552	4,181	1,873	1,364	307	1,086
Pseudo R ²	0.01	0.05	0.05	0.04	0.03	0.12	0.06	0.06	0.12	0.04	0.03	0.12	0.03

The table reports average marginal effects from a logit model where the dependent variable is an indicator of whether the observation is in the AMA dataset (i.e. has missing age at migration). Each column corresponds to different samples defined by location of birth. Each column includes all ACS observations from the listed country of birth and all AMA observations that either report the listed country of birth or do not report a country of birth but have a positive estimated probability of being born in the listed country. The full range of covariates includes 10 census divisions. Statistical significance at the 1, 5, and 10 percent levels are denoted respectively by *, **, and ***. Standard errors are in parentheses.

Table 6: AMA data—country of training and birth, when reported

LOCATION OF BIRTH	Location of Training													CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES	TOTAL
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE							
EGYPT	1,816	1	0	0	0	0	1	10	0	0	2	0	0	47	15	31	37	2	558	2,520
ETHIOPIA	2	243	0	1	0	0	0	1	0	0	1	1	0	19	3	38	3	10	232	554
GHANA	0	0	325	0	1	4	1	0	0	0	0	0	0	35	3	40	17	3	384	813
KENYA	2	0	1	104	0	0	1	0	0	2	0	4	1	28	94	11	8	167	291	714
LIBERIA	1	0	0	0	14	0	0	1	0	0	0	0	0	3	0	7	5	4	91	126
NIGERIA	1	0	0	0	0	415	1	0	1	0	0	1	0	62	9	29	8	10	342	879
OTHER EAST AFRICA	0	0	0	1	0	1	32	0	2	8	0	1	5	32	24	17	5	67	185	380
OTHER NORTH AFRICA	7	0	0	0	0	0	0	127	1	0	0	0	0	9	7	19	23	15	121	329
OTHER WEST, CENTRAL, & S'THERN AFRICA	1	0	2	0	1	7	1	0	37	5	0	0	0	49	10	51	13	4	144	325
SOUTH AFRICA	1	0	0	0	1	0	0	0	0	190	0	1	0	4	10	4	13	4	172	400
SUDAN	8	0	0	0	0	0	0	0	0	0	122	0	0	4	3	10	1	2	31	181
UGANDA	0	1	0	0	0	0	3	0	0	0	0	37	0	9	27	8	3	45	69	202
ZIMBABWE	0	0	0	0	0	0	1	0	0	25	0	0	19	0	3	2	0	1	37	88
CARIBBEAN	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	2
ENGLISH SPEAKING	6	0	3	0	0	19	4	2	0	16	0	0	3	0	0	0	0	0	0	53
EUROPE	7	1	3	0	0	11	1	3	0	4	1	0	0	0	0	0	0	0	0	31
REST OF WORLD	165	0	1	0	0	5	1	12	0	36	0	3	2	0	3	0	0	0	0	228
SOUTH ASIA	18	0	1	1	0	28	20	4	0	3	0	2	1	0	1	0	0	0	0	79
UNITED STATES	63	1	0	0	3	21	3	28	0	21	2	7	2	0	7	0	0	0	0	158
TOTAL	2,098	247	336	107	20	511	70	189	42	310	128	57	33	301	219	267	136	334	2,657	8,062

Table 7: AMA data—country of training and birth predictions

LOCATION OF BIRTH	Location of Training													CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES	TOTAL	
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE								
EGYPT	1,712	1	0	0	0	0	1	13	0	0	4	0	0	0	0	0	0	0	0	0	1,732
ETHIOPIA	2	267	0	0	0	0	0	2	0	0	0	1	0	0	0	0	0	0	0	0	272
GHANA	0	0	347	0	0	22	1	0	0	0	0	0	0	0	0	0	0	0	0	0	370
KENYA	1	0	1	65	0	0	2	0	0	8	0	2	1	0	0	0	0	0	0	0	80
LIBERIA	1	0	0	0	23	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	24
NIGERIA	1	0	0	0	0	2,061	0	0	0	0	0	1	0	0	0	0	0	0	0	0	2,063
OTHER EAST AFRICA	0	0	0	1	0	6	23	0	4	28	0	1	22	0	0	0	0	0	0	0	84
OTHER NORTH AFRICA	7	0	0	0	0	0	0	173	0	0	0	0	0	0	0	0	0	0	0	0	180
OTHER WEST, CENTRAL, & S'THERN AFRICA	1	0	2	0	4	40	3	0	50	17	0	0	0	0	0	0	0	0	0	0	116
SOUTH AFRICA	0	0	0	0	1	0	0	0	0	628	0	1	0	0	0	0	0	0	0	0	631
SUDAN	8	0	0	0	0	0	0	0	0	0	185	0	0	0	0	0	0	0	0	0	193
UGANDA	0	1	0	0	0	0	7	0	0	0	0	63	0	0	0	0	0	0	0	0	71
ZIMBABWE	0	0	0	0	0	0	0	0	0	88	0	0	33	0	0	0	0	0	0	0	122
CARIBBEAN	0	0	0	0	0	0	0	1	4	0	0	0	0	0	0	0	0	0	0	0	4
ENGLISH SPEAKING	7	0	4	0	0	108	7	3	0	53	0	0	2	0	0	0	0	0	0	0	184
EUROPE	7	1	3	0	0	123	0	3	0	12	2	0	0	0	0	0	0	0	0	0	151
REST OF WORLD	141	0	2	0	0	25	0	15	0	123	0	7	10	0	0	0	0	0	0	0	323
SOUTH ASIA	15	0	1	0	0	103	14	5	0	8	0	2	0	0	0	0	0	0	0	0	147
UNITED STATES	62	1	0	0	11	174	5	30	0	71	4	18	5	0	0	0	0	0	0	0	381
TOTAL	1,964	272	360	66	39	2,664	64	244	58	1,035	196	94	73	0	0	0	0	0	0	0	7,129

Table 8: AMA data—country of training and birth combined

LOCATION OF BIRTH	Location of Training													CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES	TOTAL
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE							
EGYPT	3,528	2	0	0	0	0	2	23	0	0	6	0	0	47	15	31	37	2	558	4,252
ETHIOPIA	4	510	0	1	0	0	0	3	0	0	1	2	0	19	3	38	3	10	232	826
GHANA	0	0	672	0	1	26	2	0	0	0	0	0	0	35	3	40	17	3	384	1,183
KENYA	3	0	2	169	0	0	3	0	0	10	0	6	2	28	94	11	8	167	291	794
LIBERIA	2	0	0	0	37	0	0	2	0	0	0	0	0	3	0	7	5	4	91	150
NIGERIA	2	0	0	0	0	2,476	1	0	1	0	0	2	0	62	9	29	8	10	342	2,942
OTHER EAST AFRICA	0	0	0	2	0	7	55	0	6	36	0	2	27	32	24	17	5	67	185	464
OTHER NORTH AFRICA	14	0	0	0	0	0	0	300	1	0	0	0	0	9	7	19	23	15	121	509
OTHER WEST, CENTRAL, & S'THERN AFRICA	2	0	4	0	5	47	4	0	87	22	0	0	0	49	10	51	13	4	144	441
SOUTH AFRICA	1	0	0	0	2	0	0	0	0	818	0	2	0	4	10	4	13	4	172	1,031
SUDAN	16	0	0	0	0	0	0	0	0	0	307	0	0	4	3	10	1	2	31	374
UGANDA	0	2	0	0	0	0	10	0	0	0	0	100	0	9	27	8	3	45	69	273
ZIMBABWE	0	0	0	0	0	0	1	0	0	113	0	0	52	0	3	2	0	1	37	210
CARIBBEAN	0	0	0	0	0	0	0	2	5	0	0	0	0	0	0	0	0	0	0	6
ENGLISH SPEAKING	13	0	7	0	0	127	11	5	0	69	0	0	5	0	0	0	0	0	0	237
EUROPE	14	2	6	0	0	134	1	6	0	16	3	0	0	0	0	0	0	0	0	182
REST OF WORLD	306	0	3	0	0	30	1	27	0	159	0	10	12	0	3	0	0	0	0	551
SOUTH ASIA	33	0	2	1	0	131	34	9	0	11	0	4	1	0	1	0	0	0	0	226
UNITED STATES	125	2	0	0	14	195	8	58	0	92	6	25	7	0	7	0	0	0	0	539
TOTAL	4,062	519	696	173	59	3,175	134	433	100	1,345	324	151	106	301	219	267	136	334	2,657	15,191

Table 9: ACS data—country of training predictions when country of birth is known

Location of Training

LOCATION OF BIRTH	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE	CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES	TOTAL
EGYPT														48	16	31	36	2	637	
ETHIOPIA														62	10	115	9	25	577	
GHANA														46	4	53	24	4	351	
KENYA														23	72	7	6	135	380	
LIBERIA														15	0	40	31	28	251	
NIGERIA														145	16	168	19	225	1,235	
OTHER EAST AFRICA														80	63	45	10	164	639	
OTHER NORTH AFRICA														18	17	28	47	32	249	
OTHER WEST, CENTRAL, & SOUTHERN AFRICA														109	20	144	35	8	381	
SOUTH AFRICA														37	76	33	113	32	1,002	
SUDAN														4	2	11	7	3	23	
UGANDA														3	34	15	2	91	81	
ZIMBABWE														0	10	6	0	2	194	
CARIBBEAN																				
ENGLISH SPEAKING																				
EUROPE																				
REST OF WORLD																				
SOUTH ASIA																				
UNITED STATES																				
TOTAL														590	341	697	339	753	6,000	

Table 10: Predictions combined for complete location of birth and education matrix

LOCATION OF BIRTH	Location of Training													CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES	TOTAL
	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST, CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE							
EGYPT	3,528	2	0	0	0	0	2	23	0	0	6	0	0	48	16	31	36	2	637	4,332
ETHIOPIA	4	510	0	1	0	0	0	3	0	0	1	2	0	62	10	115	9	25	577	1,320
GHANA	0	0	672	0	1	26	2	0	0	0	0	0	0	46	4	53	24	4	351	1,183
KENYA	3	0	2	169	0	0	3	0	0	10	0	6	2	23	72	7	6	135	380	819
LIBERIA	2	0	0	0	37	0	0	2	0	0	0	0	0	15	0	40	31	28	251	405
NIGERIA	2	0	0	0	0	2,476	1	0	1	0	0	2	0	145	16	168	19	225	1,235	4,290
OTHER EAST AFRICA	0	0	0	2	0	7	55	0	6	36	0	2	27	80	63	45	10	164	639	1,135
OTHER NORTH AFRICA	14	0	0	0	0	0	0	300	1	0	0	0	0	18	17	28	47	32	249	708
OTHER WEST, CENTRAL, & S'THERN AFRICA	2	0	4	0	5	47	4	0	87	22	0	0	0	109	20	144	35	8	381	868
SOUTH AFRICA	1	0	0	0	2	0	0	0	0	818	0	2	0	37	76	33	113	32	1,002	2,115
SUDAN	16	0	0	0	0	0	0	0	0	0	307	0	0	4	2	11	7	3	23	374
UGANDA	0	2	0	0	0	0	10	0	0	0	0	100	0	3	34	15	2	91	81	339
ZIMBABWE	0	0	0	0	0	0	1	0	0	113	0	0	52	0	10	6	0	2	194	378
CARIBBEAN	0	0	0	0	0	0	0	2	5	0	0	0	0							6
ENGLISH SPEAKING	13	0	7	0	0	127	11	5	0	69	0	0	5							237
EUROPE	14	2	6	0	0	134	1	6	0	16	3	0	0							182
REST OF WORLD	306	0	3	0	0	30	1	27	0	159	0	10	12							548
SOUTH ASIA	33	0	2	1	0	131	34	9	0	11	0	4	1							225
UNITED STATES	125	2	0	0	14	195	8	58	0	92	6	25	7							532
TOTAL	4,062	519	696	173	59	3,175	134	433	100	1,345	324	151	106	590	341	697	339	753	6,000	19,997

Table 11: Location of birth for five largest countries in terms of training

	EGYPT		ETHIOPIA		GHANA		NIGERIA		SOUTH AFRICA	
Born at home	3,528	87%	510	98%	672	97%	2,476	78%	818	61%
	(15)		(2)		(4)		(97)		(33)	
Born in other African country	44	1%	5	1%	6	1%	80	3%	181	13%
	(5)		(2)		(2)		(30)		(24)	
Born in US	125	3%	2	0%	0	0%	195	6%	92	7%
	(8)		(1)		(0)		(58)		(17)	
Born in rest of the world	365	9%	2	0%	18	3%	424	13%	254	19%
	(13)		(1)		(4)		(69)		(23)	
TOTAL TRAINED	4,062	100%	519	100%	696	100%	3,175	100%	1,345	100%

The table reports estimated totals as described in the text for selected countries. Bootstrapped standard errors are in parentheses using 100 replications.

Table 12: Location of training for five largest countries in terms of birth

	EGYPT		ETHIOPIA		GHANA		NIGERIA		SOUTH AFRICA	
Trained at home	3,528	81%	510	39%	672	57%	2,476	58%	818	39%
	(15)		(2)		(4)		(97)		(33)	
Trained in other African country	33	1%	11	1%	29	2%	6	0%	6	0%
	(6)		(3)		(22)		(3)		(6)	
Trained in US	637	15%	577	44%	351	30%	1,235	29%	1,002	47%
	(192)		(174)		(38)		(219)		(175)	
Trained in rest of the world	133	3%	222	17%	131	11%	573	13%	290	14%
	(52)		(65)		(33)		(33)		(64)	
TOTAL BORN	4,332	100%	1,320	100%	1,183	100%	4,290	100%	2,115	100%

The table reports estimated totals as described in the text for selected countries. Bootstrapped standard errors are in parentheses using 100 replications.

Table 13: Standard errors for all predictions combined

LOCATION OF BIRTH	EGYPT	ETHIOPIA	GHANA	KENYA	LIBERIA	NIGERIA	OTHER EAST AFRICA	OTHER NORTH AFRICA	OTHER WEST CENTRAL, & SOUTHERN AFRICA	SOUTH AFRICA	SUDAN	UGANDA	ZIMBABWE	CARIBBEAN	ENGLISH SPEAKING	EUROPE	REST OF WORLD	SOUTH ASIA	UNITED STATES
EGYPT	14.7	0.9	-	-	-	-	1.4	4.9	-	-	3.8	-	-	22.3	7.1	12.8	15.5	1.6	192
ETHIOPIA	1.1	2.4	-	0.4	-	-	-	1.9	-	-	0.7	1.7	-	26.4	7.3	39.8	6.7	12.2	174.4
GHANA	-	-	3.9	-	1.8	21.5	2.1	-	-	-	-	-	-	15.8	3.6	17.3	10.2	2.5	38.5
KENYA	1.2	-	1.1	1.7	-	-	4.1	-	-	6.2	-	2.2	1.8	10.3	17.8	3.5	2.8	32.4	72.8
LIBERIA	0.9	-	-	-	11.1	-	-	1.3	-	-	-	-	-	26.8	-	64.2	28.8	30.8	120.6
NIGERIA	1	-	-	-	-	97	0.8	-	-	-	-	2.4	-	48	6.4	79.1	9.2	115.6	219.4
OTHER EAST AFRICA	-	-	-	0.8	-	8.3	8.3	-	4.1	10.4	-	1.2	15.9	35.2	36.4	19.9	5.4	52.1	241.7
OTHER NORTH AFRICA	2.4	-	-	-	-	-	-	9.6	-	-	-	-	-	10.4	9	14.9	21.2	19.6	103.6
OTHER WEST, CENTRAL, & S'THERN AFRICA	0.9	-	1.8	-	4.8	23.8	3.1	-	5.4	8	-	-	-	40.2	19.3	49	13.9	7.6	108.6
SOUTH AFRICA	0.6	-	-	-	5.8	-	-	-	-	32.6	-	2.6	-	25.2	24.9	16.2	35	23.2	174.6
SUDAN	3.1	-	-	-	-	-	-	-	-	-	6.1	-	-	8.4	8.7	12.1	18.4	6.8	21.4
UGANDA	-	1.3	-	-	-	-	3.8	-	-	-	-	10.5	-	2.7	22.3	12.9	1.6	37.3	56.8
ZIMBABWE	-	-	-	-	-	-	0.8	-	-	19.4	-	-	19.2	-	6.9	10.9	-	5.6	99.1
CARIBBEAN	-	-	-	-	-	-	-	0.9	4.2	-	-	-	-	-	-	-	-	-	-
ENGLISH SPEAKING	2.8	-	2.5	-	-	33.4	5.5	2	-	14.2	-	-	5.2	-	-	-	-	-	-
EUROPE	2.5	1.3	2	-	-	61.8	1.5	3.1	-	6.4	2.5	-	-	-	-	-	-	-	-
REST OF WORLD	11.1	-	1.9	-	-	24	0.3	4.9	-	20.7	-	6.4	11.1	-	-	-	-	-	-
SOUTH ASIA	3.8	-	0.8	1.5	-	28.6	7.4	2.7	-	4.4	-	3.6	5.1	-	-	-	-	-	-
UNITED STATES	7.9	0.8	-	-	11.5	57.8	5.5	6.8	-	17.2	4	8.4	6.4	-	-	-	-	-	-

KNOMAD
Global Knowledge Partnership on Migration and Development

